

Wheat Stem Rust Genome Project: First look under the hood.

Les J. Szabo, USDA–ARS Cereal Disease Laboratory, University of Minnesota, St. Paul, MN 55108, USA.

Puccinia graminis f. sp. *tritici*, the casual agent of stem rust of wheat, has caused major epidemics over the last century. Renewed awareness of the threat to wheat production posed by this disease has occurred with the emergence of a new, virulent race (TTKS) in Eastern Africa. *P. graminis* is a dikaryotic fungus with five spore stages and two plant hosts. Whole-genome shotgun coverage (12X) has been generated by paired-end sequencing from three libraries generated by randomly sheared, total genomic libraries. A 6.8 X draft assembly containing 392 supercontigs (N50 911 kb) and covering 89 Mb was constructed. A restriction fingerprint map containing 1,969 contigs (10.8 X) was generated from approximately 22,000 fosmid clones. The restriction fingerprint map and draft sequence assembly are highly similar with 99% of the sequence assembly being placed on the fingerprint map. Annotation of the draft assembly predicted 20,567 genes, with approximately one-half of the gene calls supported by homology to sequence data in public databases and/or *P. graminis* EST database. Approximately 57,000 ESTs were generated from three cDNA libraries (urediniospores, germinated urediniospores, and teliospores), which were assembled into 6,465 synthetic contigs. Thirty-six percent of the EST contigs could be annotated using GO categories. The majority of the EST synthetic contigs were stage-specific, with only 8% common to all three libraries.

Genomic and metabolic analysis of R gene-mediated defense pathways in wheat.

Ming-Shun Chen, Plant Science and Entomology Research Unit, USDA–ARS, 4008 Throckmorton Hall, Kansas State University, 123 Waters Hall, Manhattan, KS 66506, USA.

Plant defense involves three major steps: surveillance, signal transduction, and the production of defense chemicals. A great deal of information on signal perception and transduction has been revealed in numerous plant-parasite systems. Here we investigated changes in metabolic pathways that might lead to accumulation of defense chemicals using genomic and metabolic profiling. The wheat-Hessian fly interaction was used in this study. Hessian fly is an insect but shares many features with plant pathogens, being sessile during feeding stages and having avirulence genes that match plant resistance genes in gene-for-gene relationships. Many genes involved in carbon/nitrogen metabolism were differentially regulated during compatible and incompatible interactions. During compatible interactions, the attacking site became a carbon sink. Photoassimilates were transported to the attacking site from other parts of the plant. Part of the transported photoassimilates were converted into amino acids through coordinated activation of key metabolic pathways including glycolysis, the tricarboxylic acid cycle, and amino acid synthesis pathways. In contrast, the attacking site became a nitrogen sink during incompatible interactions. Nitrogen was transported to the attacking site from other parts of the plant in the form of asparagine. The transported nitrogen was likely converted into defensive secondary metabolites. Our data suggested that the formation of a carbon sink and the conversion of C-compounds into N-compounds at the feeding site is a necessary condition for Hessian fly larvae to survive and develop in susceptible plants, whereas the formation of a nitrogen sink and the increase in phenylpropanoids and other secondary metabolites may be part of the resistance mechanism.

Fine mapping of the Lr46/Yr29 locus in wheat.

Leandro Perugini ¹, Maria Mateos-Hernandez ², John Fellers ³, Qijian Song ⁴, Perry Cregan ⁴, Scot Hulbert ⁶, Robert Bowden ³, Ravi Singh ², and Gina Brown-Guedira ^{1,5}.

¹ Department of Crop Science, North Carolina State University, Raleigh, NC 27695, USA; ² CIMMYT, International, Apdo Postal 6-641, Mexico City, Mexico 06600; ³ USDA–ARS and Department of Plant Pathology, Kansas State University, Manhattan, KS 66506, USA; ⁴ USDA–ARS, Beltsville Agricultural Research Center, Beltsville, MD 20705, USA; ⁵ USDA–ARS and Department of Crop Science, North Carolina State University, Raleigh, NC 27695, USA; and ⁶ Department of Plant Pathology, Washington State University, Pullman, WA 99164, USA.

Rust diseases, including leaf rust (*Puccinia triticina*), stripe rust (*P. striiformis* f. sp. *tritici*), and stem rust (*P. graminis* f. sp. *tritici*) are globally the most important diseases of wheat. Genetic resistance is the best form of control but frequent race changes typically render new varieties susceptible within a few years after release. For typical resistance genes, the

mechanism of resistance and the basis of race specificity are due to gene-for-gene interactions. However, some disease resistance genes are thought to be durable, because they are not dependent on the recognition of a single *Avr* gene product from the pathogen by an *R* gene. The adult-plant resistance gene *Lr46* has provided non-race-specific resistance to leaf rust that has remained effective for more than 30 years. The gene also has a pleiotropic effect on resistance to stripe rust (*Yr29*). Using recombinant, chromosome substitution line populations, we previously located *Lr46* on the terminal region of the long arm of chromosome 1B that is syntenic to chromosome 5L of rice. To fine map the *Lr46* gene region, high resolution mapping (HRM) populations were developed that represent 4,100 gametes. The EST-derived, STS marker *XSTS3680* that co-segregated with *Lr46* in the original mapping populations was mapped 0.15 cM distal to *Lr46* in the HRM populations. A BAC contig of the *Lr46* region is being constructed. New SSR and SNP markers identified from the BAC clones and linked to *Lr46* have been evaluated on a set of diverse wheat lines to determine their usefulness for marker-assisted selection. Gene expression studies can complement map-based cloning efforts, because expression data can be used for identifying candidate genes, identifying expression markers, and for generating and testing hypotheses about genetic resistance mechanisms. To identify transcripts associated with *Lr46*-mediated adult-plant resistance, the Affymetrix Wheat GeneChip Microarray was used to identify transcriptional changes in isogenic lines with and without *Lr46*. Considering the increasing worldwide use of *Lr46* and other adult-plant genes for durable rust resistance, it is essential to obtain a greater understanding of their mechanisms of resistance. Also essential is obtaining the best possible markers for breeding for durable resistance.

Genetic analysis of host-toxin interactions in the wheat–Stagonospora nodorum pathosystem.

Justin D. Faris¹, Timothy L. Friesen¹, Huangjun Lu², Leela Reddy³, Zencui Zhang³, Zhaohui Liu², Steven S. Xu¹, Steven W. Meinhard², John P. Fellers⁴, Richard P. Oliver⁵, Peter S. Solomon⁵, Shiaoman Chao¹, Chenggen Chu³, Nilwala Abeysekara³, and Steven R. Scofield⁶.

¹ USDA–ARS Cereal Crops Unit, Northern Crop Science Laboratory, Fargo, ND 58105, USA; ² Department of Plant Pathology, North Dakota State University, Fargo, ND 58105, USA; ³ Department of Plant Sciences, North Dakota State University, Fargo, ND 58105, USA; ⁴ USDA–ARS Plant Science and Entomology Research Unit, Kansas State University, Manhattan, KS 66506, USA; ⁵ Australian Centre for Necrotrophic Fungal Pathogens, Western Australian State Agricultural Biotechnology Centre, Murdoch University, Western Australia 6150; and ⁶ USDA–ARS Crop Production and Pest Control Research Unit, Purdue University, West Lafayette, IN 47907, USA.

Stagonospora nodorum, causal agent of *Stagonospora nodorum* blotch (SNB) in wheat, produces multiple necrosis-inducing host-selective toxins (HSTs) that interact with dominant host sensitivity genes to cause disease. Absence of either the toxin or the dominant host gene precludes recognition and results in an incompatible (resistant) response. Therefore, these host-toxin interactions are mirror images of classical gene-for-gene interactions. One of the first HSTs identified in this system was SnToxA, which was horizontally transferred from *S. nodorum* to the tan spot pathogen *Pyrenophora tritici-repentis* around 1941. This event is considered to have been significant for the establishment of tan spot as a pathogen. Sensitivity to SnToxA is governed by the *Tsn1* gene on the long arm of chromosome 5B. To date, eight additional toxins, designated SnTox1 through SnTox8, have been identified, and their corresponding host sensitivity genes, designated *Snn1* through *Snn8*, have been mapped to wheat chromosome arms 1BS, 2DS, 5BS, 4BL, 5BS, 6AL, 5DS, and 3DL, respectively. Genetic analysis of several host-toxin interactions indicates that they play important roles in the development of disease in adult plants as well as seedlings, and their effects are mostly additive. To gain a better understanding of compatible host-toxin interactions at the molecular level, we have embarked on the positional cloning of two host-sensitivity genes: *Tsn1* on 5BL and *Snn1* on 1BS. Toward the map-based cloning of *Tsn1* on chromosome 5B, we sequenced and assembled chromosome 5A and 5B BAC contigs spanning the gene. Evaluation of gene content and micro-colinearity between the orthologous regions of 5A, 5B, and rice chromosome 9 indicated the 5A region and rice share a higher level of micro-colinearity than the 5B region does with rice due to the presence of numerous transpositions, deletions, and rearrangements present in the wheat 5B region. In addition, the 5B *Tsn1* candidate region is nearly 4 times larger than the corresponding region of 5A due to the presence of additional genes and transposable elements. At least ten genes exist within the 350 kb *Tsn1* candidate-gene region, and they are currently being validated by comparative sequence analysis of *Tsn1*-disrupted mutants and virus-induced gene silencing. An important applied by-product of this research is the development of efficient PCR-based markers for *Tsn1*, which are being used to introgress SnToxA insensitivity into adapted germ plasm. Overall, this research demonstrates the potential of the wheat–*S. nodorum* pathosystem to be an excellent toxin-based inverse gene-for-gene model.