

of marker systems will yield a better contiguous map. The mapping resolution of these RH panels estimated on the basis of markers spanning known distances was <140 kb. Two sets of informative lines carrying breaks in multiple D-genome chromosomes were selected from *Ae. tauschii* DGRH1s (399 lines) and Chinese Spring DGRH1s (300 lines). First generation RH maps based on 178 lines and 676 markers (641 DaRT and 35 SSR) showed a 17:1 map ratio cR/cM when compared with the genetic maps. A NimbleGen array has been designed and tested for high-throughput mapping, and a total of ~30,000 retro-junction markers and ~6,000 gene-based markers, specific to the D genome were identified. The selected DGRH1 lines currently are undergoing genotyping with this array and, once analyzed, will provide a very dense scaffold for the assembly of the D genome of wheat. This research also provides valuable resources for fine mapping and map based cloning studies of genes present on the D genome along with an unprecedented view into the evolution of grass genomes (<http://avena.pw.usda.gov/RHmapping/>).

### ***Poster 8. Construction of a radiation hybrid map for chromosome 6B of common wheat.***

Shyota Watanabe, Miyuki Nitta, Takashi R. Endo, and Shuhei Nasuda. Laboratory of Plant Genetics, Graduate School of Agriculture, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan.

It is well known that recombination events in wheat are not evenly distributed along the length of chromosomes. This is the case with chromosome 6B, which is the target chromosome assigned to Japan by the International Wheat Genome Sequencing Consortium. The genetic maps of chromosome 6B so far constructed are heavily populated by markers in the pericentromeric region and scarcely in the telomeric regions. This represents that recombinations mostly take place in the telomeric ends of the chromosome. Therefore, we would face problems in determining orientation of contigs in the pericentromeric region if we solely depend on the genetic maps. Our objective of the current study is to establish a radiation hybrid (RH) mapping panel that can be useful in determining marker orders in the recombination-poor, pericentromeric region of chromosome 6B. We crossed nullisomic 6B-tetrasomic 6A plants of Chinese Spring (CS) wheat with the pollen freshly irradiated by  $\gamma$ -ray (15 Gy). We sowed 2,171 M0 seeds and extracted genomic DNA from 461 (21.2%) surviving plants. Additionally, we used 12 6B deletion lines (five deletions in 6BS and seven deletions in 6BL; obtained from NBRP-Wheat, Japan). We scored the presence or absence of 21 previously reported 6B-specific SSR markers and four newly developed EST-based markers. We analyzed the data by CarthaGene software to construct a RH map with the default setting. The resulting RH map consisted of two linkage groups, corresponding to the short and long arms. The gap between the linkage groups may be due to the absence of markers in a pericentromeric bin (6BS-CEN-0.25). The marker order is consistent with that of bin mapping and largely with a previously reported genetic map. All the RH-map markers occupied individual loci. Two pairs of markers genetically mapped to the same loci in the pericentromeric region of 6B were completely separated from each other in the RH map. This result indicates that RH mapping with our panel has better resolution in proximal region of 6B.

### ***Poster 9. A NimbleGen comparative genomic hybridization array for high-throughput physical mapping of genome specific repeat junction and gene-based markers in the D genome of hexaploid wheat.***

Thomas Drader<sup>1</sup>, Lingli Dong<sup>1,4</sup>, Yi Wang<sup>1,4</sup>, Ajay Kumar<sup>2</sup>, Vijay K. Tiwari<sup>3</sup>, Muhammad Iqbal<sup>2</sup>, Jeff M. Leonard<sup>3</sup>, Gerard Lazo<sup>1</sup>, Anne Denton<sup>2</sup>, Shahryar Kianian<sup>2</sup>, Ming-Cheng Luo<sup>4</sup>, and Yong Gu<sup>1</sup>.

<sup>1</sup> USDA-ARS-WRRC, Albany, CA 94710, USA; <sup>2</sup> North Dakota State University, Fargo, ND 58105, USA; <sup>3</sup> Oregon State University, Corvallis, OR 97370, USA; and <sup>4</sup> University of California, Davis, CA 95616, USA.

Mapping and map-based cloning of agriculturally and economically important traits remain a great challenge in complex highly repetitive genomes such as the grass tribe, Triticeae. This limitation is primarily based upon the availability of polymorphic markers and frequency of genetic recombination. Most markers are gene-based, derived from polymorphisms within coding regions. Non-gene-based markers, such as repeat junction markers, are derived from the noncoding intergenic space. Repeat junction markers take advantage of the repetitive nature of the wheat genome, providing random and equal distribution of these markers throughout the genome and can facilitate mapping efforts. Repeat junction markers are designed upon the junction of nested repetitive elements, and approximately 90% of these markers have been determined to behave as a single copy locus and are genome specific. Repeat junction markers were designed from 454 genome sequences of the wheat D-genome progenitor, *Aegilops tauschii*. Mapping of *Ae. tauschii* repeat junction