

Variations in patterns of gene expression are central to evolution. A large-scale screen of barley germplasm has identified an alternative de-repressor of lodicule development SV235. The relevant *HvAP2* coding sequence is identical to that of *cly1.b*, so that *HvAP2* down-regulation is not associated with miR172-directed degradation; instead, transcriptional repression appears to be induced by the maintenance of the *HvAP2* promoter in a hyper-methylated state during the period when the vascular tissue in the lodicule would normally develop. The cleistogamous type can also arise in the presence of non-cleistogamous type lodicule development. In certain *Cly1.a* x *cly1.b* hybrids, the spikes remain within the boot during anthesis, and the lodicules remain shrunken even after the emergence of the spike from the boot.

Genetics and molecular evolution of a 3.1-Mb genomic region harboring both wheat prolamin and disease resistance gene families.

Lingli Dong ^{1,2}, Naxin Huo ^{1,2}, Yi Wang ^{1,2}, Karin Deal ², Frank You ³, Jan Dvorak ², Olin D. Anderson ¹, Ming-Cheng Luo ², and **Yong Q. Gu** ¹.

¹ United States Department of Agriculture-Agricultural Research Service, Western Regional Research Center, Albany, CA 94710, USA; ² Department of Plant Sciences, University of California, Davis, CA 95616, USA; and ³ Cereal Research Center, Agriculture and Agri-Food Canada, 195 Dafoe Road, Winnipeg, Manitoba, Canada, R3T 2M9.

Several important wheat prolamin and disease resistance gene loci have been mapped at similar locations in the short arm of the group-1 chromosomes in wheat. However, genetics and structural organization of genomic regions harboring these important traits have not been well characterized. We therefore sequenced a 3.1-Mb region harboring prolamin multigene families in the genome of *Ae. tauschii*, the D-genome donor of hexaploid bread wheat. The sequence revealed a much higher gene density (98 genes; one gene/31 kb) in the sequenced region than the average value (ca. one gene/112 kb) expected for the wheat D genome. The high gene density is primarily due to the large number of duplicated genes, present either in tandem or interspersed with other genes. In addition to different types of prolamin gene families (γ -gliadin, ω -gliadin, and LMW-glutenin), multiple NBS-LRR disease resistance gene homologues of *Lr21* (resistance to leaf rust) and *Pm3* (resistance to powdery mildew) were identified. Furthermore, leucine-rich receptor protein kinase (LRK) genes, representing a different class of resistance genes, also were highly duplicated in this region. Comparative analyses indicated that the orthologous regions in the rice, *Brachypodium*, and sorghum genomes are highly conserved, whereas in *Ae. tauschii*, only 16 out of 98 genes are syntenic. Most of these highly duplicated genes are unique in the *Ae. tauschii* genome, suggesting rapid evolution in this region. Genetic analysis revealed the sequenced region spanned over 10-cM genetic distance with 13 markers mapped to this region. Recombination rates ranged from ca. 200 to 1,000 kb/cM between two marker intervals. Both co-evolution and independent evolution in different gene families were observed. Mechanisms underlying the molecular evolution of prolamin and resistance gene families in this complex region will be presented.

SESSION V: YOUNG TRITICEAE RESEARCHERS

Genetic provenance and genetic providence in a diverse crop.

Ana M. Gonzales and Peter L. Morrell. Department of Agronomy and Plant Genetics, University of Minnesota, St. Paul, MN, USA.

Multiple factors converge to contribute to high levels of genetic and phenotypic diversity in barley. The progenitor species is ecotopically and genetically diverse, with a large geographic range and large effective population size. Cultivated barley arose from multiple source populations and became adapted to diverse agronomic conditions across much of Eurasia and North Africa over several millennia of human prehistory. Exploiting this diverse adaptive history for barley improvement is challenging if methods are limited to top-down approaches based on genotype-phenotype associations. A new generation of bottom-up approaches provides the opportunity to investigate the genetic differentiation and genetic provenance of adaptive allelic variation and associations between SNPs and ecogeographic variables. Both sets of approaches provide new opportunities for associating genetic variation with phenotypic traits of agronomic importance, including disease resistance and climate adaptation. We make use of genotyping data from 2,417 worldwide barley accessions from the USDA National Small Grains Collection (landraces and cultivars) genotyped with 7,800 SNPs.