

ITEMS FROM THE RUSSIAN FEDERATION

AGRICULTURAL RESEARCH INSTITUTE FOR THE SOUTH-EAST REGIONS
Department of Genetics, Laboratory of Genetics and Cytology, 7 Toulaikov St., Saratov,
410010, Russian Federation.

The influence of a translocation with the Lr19+Lr26 combination on grain productivity and bread-making quality in spring bread wheat.

S.N. Sibikeev, A.E. Druzhin, and S.A. Voronina.

At the Agricultural Research Institute for the South-East Regions (ARISER), NILs based on the Saratov-bred spring bread wheat cultivar L503 and carrying translocations with the *Lr19+Lr26* combination, prospective lines were produced and studied. The data from 2005–11 indicate that the interaction of these translocations positively influences grain yield (Table 1). During this period, leaf rust epidemics were observed three times (2005, 2006, and 2008) and drought conditions four times (2007, 2009, 2010, and 2011). Grain productivity significantly increased in two leaf rust epidemics in lines with *Lr19+Lr26*, and twice under drought conditions. In the hard drought of 2010, spring bread wheat plants died. The main limiting factor for the use of T1BL·1RS translocations in wheat breeding is the influence on bread-making quality. The decrease of bread-making quality is coupled with genes for disease resistance and *Sec1*. In the NILs produced in the genetic background of cultivar L503, the presence of *Lr19+Lr26* translocations did not influence the grain protein content or gluten values. The increase in grain yield of lines containing *Lr19+Lr26*-translocations was not accompanied by decreased of grain protein content. Dough extensibility (P) and strength of flour (W) were not significantly lower in the NILs and perspective lines with *Lr19+Lr26* translocations compared with L503. Similar results were obtained for bread-making qualities. Loaf volume and porosity also were not significantly lower than L503 in the NILs and lines with *Lr19+Lr26* translocations (Table 2). Thus, the *Lr19+Lr26* translocations had positive effects on resistance to disease and drought and grain productivity with good bread-making quality.

Table 1. Grain productivity, grain protein content, and gluten deformation index of L503 NILs and perspective lines of spring bread wheat with *Lr19+Lr26*.

NIL / line	Grain yield (kg/ha)	Grain protein content (%)	Gluten value	
			content (%)	strength
L503 (<i>Lr19</i>)	1,755 a	17.78	42.17	81
L503 (<i>Lr19+Lr26</i>)	2,096 bc	17.94	40.53	78
L503(<i>Lr19+Lr26</i>)/L505	2,301 c	17.79	39.32	81
LSD	47.79	NS	NS	NS

Table 2. Bread-making qualities of L503 NILs and perspective lines of spring bread wheat with *Lr19+Lr26* (dough extensibility = P and strength of flour = W).

NIL / line	Physical traits of dough			Bread-making qualities		
	P	P/L	W	loaf volume (cm ³)	porosity	crumb color
L503 (<i>Lr19</i>)	93.4	1.5	192	788	4.5	yellow
L503 (<i>Lr19+Lr26</i>)	84.6	1.4	166	760	4.4	yellow
L503(<i>Lr19+Lr26</i>)/L505	89.3	1.3	185	750	4.3	yellow
LSD	NS	NS	NS	NS	NS	

The agronomic performance of Lr19+Lr37 translocations in the set of NILs in the genetic background of the spring bread wheat cultivar Dobrynya.

S.N. Sibikeev and A.E. Druzhin .

We used a set of NILs in the genetic background of the spring bread wheat cultivar Dobrynya to associate positive agronomical traits of *Lr19* and *Lr37* translocations. The presence of both translocations in the NILs was confirmed with molecular markers; STS marker Gb Lr19 for the *Lr19* translocation and primers VENTRIUP and LN2 for the *Lr37* translocation (Gulyaeva EI et al. 2012). In the droughty conditions of 2011, NILs with *Lr19+Lr37* translocations were significant higher than Dobrynya for grain productivity. The NILs also surpassed Dobrynya in grain protein content, however, the results for gluten content and gluten strength were ambiguous. One line was higher than that of Dobrynya, and another line was equal to Dobrynya (Table 3). For dough extensibility (P) and strength of flour (W), NIL 118 was equal to and NIL 113 exceeded those of Dobrynya. Loaf volume and porosity were lower in the NILs with *Lr19+Lr37* translocations than in Dobrynya (Table 4). We conclude that translocations with the *Lr19+Lr37* combination improve drought resistance but do not depress bread-making properties.

Table 3. Grain productivity, grain protein content, and gluten deformation index of Dobrynya NILs and perspective lines of spring bread wheat with *Lr19+Lr37*.

NIL / line	Grain yield (kg/ha)	Grain protein content (%)	Gluten value	
			content (%)	strength
Dobrynya (<i>Lr19</i>)	2,695 a	14.98	37.4	80
NIL 113 Dobrynya (<i>Lr19+Lr37</i>)	2,864 b	15.75	35.4	68
NIL 118 Dobrynya (<i>Lr19+Lr37</i>)	3,156 c	16.10	40.0	78
LSD	153			

Table 4. Bread-making qualities of Dobrynya NILs and perspective lines of spring bread wheat with *Lr19+Lr37* (dough extensibility = P and strength of flour = W).

NIL / line	Physical traits of dough			Bread-making qualities		
	P	P/L	W	loaf volume (cm ³)	porosity	crumb color
Dobrynya (<i>Lr19</i>)	108	1.9	249	810	5.0	yellow
NIL 113 Dobrynya (<i>Lr19+Lr37</i>)	156	2.7	366	700	4.5	yellow
NIL 118 Dobrynya (<i>Lr19+Lr37</i>)	110	1.7	249	710	5.0	yellow

Resistance screening to a local population of leaf rust among species and relatives of bread wheat.

S.N. Sibikeev, A.E. Druzhin, T.D. Golubeva, and T.V. Kalintseva.

Screening for resistance to a local population of leaf rust among 25 species and relatives of bread wheat gave the following results: 14 lines from 10 species/relatives of bread wheat and one triticale cultivar, Satu, were resistant. Temperature sensitivity for resistance to leaf rust was found in *T. turgidum* and *T. petropavloskyi* in 2006. With the purpose of enlarging the gene pool of bread wheat for *Lr* genes, hybrids were obtained between the spring bread wheat cultivars Saratovskaya 68, Saratovskaya 70, Saratovskaya 73, and Saratovskaya 74 and lines of the resistant species.

Evaluating spring bread wheat cultivars for resistance to tan spot (Pyrenophora tritici-repentis (Died.) Drechsler.) in 2011.

S.N. Sibikeev, A.E. Druzhin, T.D. Golubeva, and T.V. Kalintseva.

Tan spot, caused by the ascomycete fungus *Pyrenophora tritici-repentis*, is one of the most serious foliar diseases of wheat. The average yield loss caused by this pathogen is 5–10%, but in epidemic years, losses can be up to 30%.

Moreover, tan spot also causes significant losses in grain quality by grain shriveling, red smudge, and black point. In recent years, this disease more often appear in the plantings of spring bread wheat in the Lower Volga Region of the Russian Federation. To screen for resistance to this pathogen, we evaluated 140 introgression lines of spring bread wheat in a nursery. Among the lines, 35 were resistant (% infection = 0) to the local tan spot population. NIL L400R, with a *Thinopyrum intermedium* 6Agi (6D) substitution was more resistant to tan spot than its sib lines L400S that lack 6Agi (Table 5). The combination of translocations T7DS·7DL-7Ae#1L (*T. elongatum*) and T6BS·6BL-6U#1L (*Ae. umbellulata*) and T7DS·7DL-7Ae#1L and 4BS·4BL-2R#1L (*S. cereale*) increase the resistance of the spring bread wheat lines to the tan spot, however, the combination T7DS·7DL-7Ae#1L and T1BL·1R#1S significantly increased susceptibility. NILs with a combination of T7DS·7DL-7Ae#1L and T3DS·3DL-3Ae#1L (*T. elongatum*) did not significantly differ on the extent of damage with respect to the cultivar (Table 5).

Table 5. The reaction of cultivars and NILs of spring bread wheat to *Pyrenophora tritici-repentis*.

Cultivar / NIL	Translocation	% infection
L 400 R	6Agi (6D)	5
L 400 S	—	30
L 503	T7DS·7DL-7Ae#1L	0
L 503 (<i>Lr26+Lr19</i>)	T7DS·7DL-7Ae#1L	0
L 503 (<i>Lr26+Lr19</i>)	T7DS·7DL-7Ae#1L + T1BL·1R#1S	5–10
L 505//L503 (<i>Lr26+Lr19</i>)	T7DS·7DL-7Ae#1L + T1BL·1R#1S	5
Dobrynya	T7DS·7DL-7Ae#1L	30
Dobrynya (<i>Lr19+Lr9</i>)	T7DS·7DL-7Ae#1L + T6BS·6BL-6U#1L	3
Dobrynya (<i>Lr19+Lr24</i>)	T7DS·7DL-7Ae#1L + T3DS·3DL-3Ae#1L	30
Dobrynya (<i>Lr19+Lr25</i>)	T7DS·7DL-7Ae#1L + 4BS·4BL-2R#1L	10

Department of Biotechnology, Laboratory of Cell Breeding, 7 Tulaikov Street, Saratov, 410010, Russian Federation.

The microclonal propagation of sterile triticales plants.

V.N. Akinina, T.I. Dyatchouk, and A.V. Pominov.

Applying biotechnology to crop improvement induces nonconventional plant breeding methodology. In biotechnology and plant breeding, preserving and propagating sterile plants, undoubled haploids, and remote hybrids for future use is important. The most frequent explant sources for morphogenetic callus induction in cereals are mature and immature embryos. To establish tissue culture systems in sterile plants, it is necessary to select the explant source.

Wheat–triticales hybrids (three genotypes) and haploids (nine genotypes) were grown in a phytootron. Somatic embryogenesis was used to produce the largest number of plantlets. Segments of immature inflorescence 2–3 cm and completely covered by leaves served as explants to establish callus cultures. A solid MS nutrient medium, supplemented with 2 mg/L 2,4-D and 2% sucrose was optimum for callus induction. Callus cultures were grown in the dark, with subculturing on fresh culture medium every 3–4 weeks. The subsequent elimination of 2,4-D from the culture medium and the addition of IAA (1 mg/L) allowed for embryogenesis. Centers of morphogenesis were milky white, compact, and modular. The start of regeneration was observed within 7–10 days. The frequency of callus induction ranged from 76.5 to 84.0% and the frequency of morphogenetic callus was from 39.1 to 46.6%.

The microclonal propagation of sterile triticales sterile was elaborated using segments of immature inflorescence as source of totipotent cultures. Embryogenic callus and plant regeneration were formed. This technology allows one to obtain clones of each genotype in an unlimited amount for use in breeding work.

Identification of leaf rust resistance genes of lines of soft spring and winter wheats.

T.S. Markelova and O.V. Ivanova.

Leaf rust resistance genes were identified in 14 introgressed lines of soft wheat carrying noncompensating translocations from *Aegilops speltoides*, *Triticum turgidum* subsp. *dicoccum* and *persicum*, and wheat cultivars from the collection of the All-Russian Institute of Plant Breeding using phytopathological and molecular methods.

After molecular screening of wheat cultivars, we discovered three with gene *Lr9* and one with *Lr19* (Table 6, p. 197). The identification of *Lr9* used SCAR marker SCS5 (Gupta et al. 2005). A characteristic 550-bp fragment was found among breeding lines 6 and 7, Clez Alta, and a control line Tc *Lr9*. The results from the molecular screening of these species correspond with the phytopathological test. Gene *Lr9* in line 11 is unidentified despite the fact that it was used when the line was created. In addition to leaf rust resistance, this line also has mildew resistance from *T. turgidum* subsp. *persicum*.

Table 6. Leaf rust (*Lr*) genes identified using DNA markers.

#	Description	<i>Lr9</i>	<i>Lr10</i>	<i>Lr19</i>	<i>Lr20</i>	<i>Lr26</i> <i>iag95</i>	<i>Lr26</i> <i>SCM9</i>	<i>Lr34</i>	<i>Lr35</i>	<i>Lr37</i>
1	Saratovskaya 29/ <i>T. persicum</i> // <i>T. dicoccum</i>				+				+	
2	Saratovskaya 55// <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
3	Saratovskaya 55// <i>T. dicoccum</i> / <i>Ae. speltoides</i>		+						+	
4	Saratovskaya 29// <i>T. persicum</i> /3/ <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
5	Saratovskaya 29// <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
6	Saratovskaya 29/ <i>T. persicum</i> // <i>Lr9</i>	+								
7	Saratovskaya 29/ <i>T. persicum</i> / <i>Lr9</i>	+								
8	Saratovskaya 29/ <i>T. persicum</i> // <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
9	Saratovskaya 55// <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
10	Saratovskaya 55// <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
11	Saratovskaya 29/ <i>T. persicum</i> // <i>Lr9</i>								+	
12	Saratovskaya 29/ <i>T. persicum</i> // <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
13	Saratovskaya 55// <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
14	Saratovskaya 29/ <i>T. persicum</i> // <i>T. dicoccum</i> / <i>Ae. speltoides</i>								+	
15	Clez Alta	+								
16	K-34612			+						
17	W464//VEE/KOEL/3/PEG//MRL/BUC		+			+	+	+		
18	CROC-1/ <i>Ae. tauschii</i> (205)//KAUZ/3/ATTILA					+	+			

A characteristic 130-bp fragment was found in sample 16 with the help of STS marker Gb *Lr19* (Prins et al. 2001). A 487-bp fragment was discovered in lines 1–5 and 8–14 with the help of marker Sr39#22 for gene *Lr35*. Gene *Lr35* is located on the short arm of chromosome 2B and is situated in the same translocation with seedling gene *Sr39*, which defends against *Puccinia graminis* race Ug99 (TTKS) (Singh et al. 2008). Diploid *Ae. speltoides* is the source of this translocation.

When marker PS10 for gene *Lr47* from *Ae. speltoides* was being applied its carriers were not discovered. Although gene *Lr35* has the potential for leaf rust resistance, examples of its application have not been given in the literature to date (McIntosh et al. 1995; Mago et al. 2009; Serfling et al. 2011). A good example of introgressing *Lr/Sr* genes from *Ae. speltoides* in Russia is the set of amphidiploid lines ((*T. dicoccum* x *Ae. speltoides*)*5//Saratovskaya 29) created at the ARJPB (Odintsova et al. 1991). These lines were widely used in many selection centres in the USSR, including SRJA of South-East. Thus, the existence of the given translocation in lines 1–5 and 8–14 is logical. Their resistance to

race *P. graminis* Ug99+*Lr24* (TTKST) (Sibikeev et al. 2011) is important to characteristics of these lines. Leaf and stem rust resistance, possibly caused by *Lr* and *Sr* cohesion, supports the existence of *Lr35/Sr39* genes in lines 1–5 and 8–14.

Although *Lr10* was identified with the primers F 12245 and *Lr10*-6/r2 (Schachermayr et al. 1997), an amplicon with a molecular weight of 310 bp was discovered in lines 3 and 17.

Using the primers STS638-L and STS638-R for gene *Lr20* (Neu et al. 2002), a marker of 540 bp was found in line 1 and in the control Thatcher with *Lr20*; line 1 includes wild *T. persicum* and *T. dicoccum* in the pedigree. Today, *Lr20* has lost its efficiency practically everywhere (McIntosh et al. 1995; Khan et al. 2005).

To identify gene *Lr26*, two markers, SCM9 (Mago et al. 2002) and iag95 (Weng et al. 2007), were used. A characteristic 207-bp component of SCM9 and an ~1,000-bp iag95 marker were identified in lines 17 and 18, which corresponds to their gender lines.

By applying phytopathological and molecular methods and also analyzing pedigrees, we discovered *Lr* genes and their combinations in 14 introgressive lines and in four soft spring wheats. These results are very important for programs that use the application of certain *Lr* genes and their combinations in practical selection.

References.

- Gupta SK, Charpe A, Koul S, et al. 2005. Development and validation of molecular markers linked to an *Aegilops umbellulata*-derived leaf rust- resistance gene, *Lr9*, for marker-assisted selection in bread wheat. *Genome* 48(5):823-830.
- Khan RR, Bariana HS, Dholakia BB, et al. 2005. Molecular mapping of stem and leaf rust resistance in wheat. *Theor Appl Genet* 111(5):846-850.
- Mago R, Spielmeyer W, Lawrence G, et al. 2002. Identification and mapping of molecular markers linked to rust resistance genes located on chromosome 1RS of rye using wheat-rye translocation lines. *Theor Appl Genet* 104:131-134.
- Mago R, Zhang P, Bariana HS, et al. 2009. Development of wheat lines carrying stem rust resistance gene *Sr39* with reduced *Aegilops speltoides* chromatin and simple PCR markers for marker-assisted selection. *Theor Appl Genet* 124:65-70.
- McIntosh RA, Wellings CR, and Park RF. 1995. Wheat rusts: an atlas of resistance genes. CSIRO, Australia, and Kluwer Academic Publishers, Dordrecht, the Netherlands. 200 pp.
- Odintsova IG, Agafonova NA, and Boguslavskiy RL. 1991. The spring bread wheat introgression lines, carrying resistance to leaf rust transferred from *Aegilops speltoides*. In: *Proc Appl Bot, Genet and Breed, Vavilov Inst Res* 142:106-110.
- Prins R, Groenewald J, Marais G, et al. 2001. AFLP and STS tagging of *Lr19*, a gene conferring resistance to leaf rust in wheat. *Theor Appl Genet* 103(6):618-624.
- Schachermayr G, Feuillet C, and Keller B. 1997. Molecular markers for the detection of the wheat leaf rust resistance gene *Lr10* in diverse genetic backgrounds. *Mol Breed* 3:65-74.
- Serfling A, Krämer I, Lind V, et al. 2011. Diagnostic value of molecular markers for *Lr* genes and characterization of leaf rust resistance of German winter wheat cultivars with regard to the stability of vertical resistance. *Eur J Plant Path* 130(4):559-575.
- Sibikeev SN, Markelova TS, Druzhin AE, Vedeneeva ML, and Singh D. 2011. Evaluation of the introgression spring bread wheat lines of Agricultural Research Institute for South-East Regions bred for resistance to race of *Puccinia graminis* f. sp. *tritici* Ug99 + *Sr24* (TTKST). *Rus Agric Sci* 2:3-5.
- Singh RP, Hodson DP, Huerta-Espino J, et al. 2008. Will stem rust destroy the world's wheat crop? *Adv Agron* 98:310.
- Weng Y, Azhaguvel P, Devkota RN, and Rudd JC. 2007. PCR-based markers for detection of different sources 1AL·1RS and 1BL·1RS wheat-rye translocations in wheat background. *Plant Breed* 126:482-486.

Laboratory of Winter Bread Wheat Breeding, 7 Toulaiikova St., Saratov, 410010,
Russian Federation.

The winter wheat cultivar Kalach 60.

J. Svistunov, A. Pryanichnicov, A. Zavorotina, A. Sergeeva, V. Uvarova, and N. Larionova.

Breeding of the winter wheat cultivars at ARISER began in 1915. The main task in the first stage of selection was creating cultivars adapted to the steppe zone of the Volga Region. Since the beginning, 26 cultivars have been developed from which seven are currently patented and registered.

Kalach 60 is an intensive type cultivar; to obtain high yield and high-quality grain, it requires the application of fertilizers and observing agronomical practices. Kalach 60 was developed using individual selection. The cultivar is of the variety lutescens. Plant height is 20–30 cm shorter compared to that of Mironovskaya 808. The straw is thick. Kalach 60 has increased winter hardiness, lodging and drought resistance, early mature growth, high grain quality, and high grain yield (2.7 t/ha). These characters allow the use of Kalach 60 in intensive farming to obtain high-quality grain. Early ripening allows Kalach 60 to be used in a system used by large farms, which sow winter crops on large areas.

In 2011, Kalach 60 was planted in the Saratov region on 13.7×10^3 hectares (2.8%). In 2012, it was listed in the National Register of the Russian Federation in the lower Volga Region.

RESEARCH DEPARTMENT OF “AGROPARK”

Gostynichnaya, 4B Moscow 127106, Russian Federation.

The role aluminum salts ($AlCl_3$, $Al(NO_3)_3$, and $Al_2(SO_4)_3$) in the formation of spring wheat resistance to aluminum toxicity.

N.V. Poukhalskaya, Y.R. Ziangirova, and F.D. Davadov, and N.I. Pavlova (Pryanishnikov All Russian Research Institute of Agriculture and Solid Science, Pryanishnikova, 31. Moscow 127550, Russian Federation) and S.L. Ignatyeva (Russian State Agrarian University – MTAA named after K.S. Timiryazev).

We are researching the toxicity of aluminum, at present, because of insufficient data about the toxicity of various Al^{+3} salts (sulfates, nitrates, and chlorides). Our data show that aluminum toxicity changes in the presence of different ions. We studied the comparative degree of toxicity of aluminum ions when using sulfates, nitrates, and chlorides on spring wheat.

Materials and methods. Aluminum tolerance was observed at germination in laboratory experiments under controlled environmental conditions. Seedlings of two cultivars of spring wheat were grown under greenhouse conditions. Sprouts of plants were grown on solutions of $AlCl_3$, $Al(NO_3)_3$, and $Al_2(SO_4)_3$ with various concentration of aluminum (0, 0.4, 3, 5.13, 15, and 40 mg of Al/L. Absorption from the solutions was measured in vessels each containing the same number of plants. Plants were grown on plastic floats (rafts) on the surface of the solution. The seed was planted in apertures in the floats. Experiments also were made in rolled culture and in soil. The same aluminum salts were used at 1.3 mg/100 g soil for the vegetative trials.

Results and discussion. During germination, wheat plants reduce the absorption from solutions in different ways. Absorption decreases as follows: $Al(NO_3)_3 = Al_2(SO_4)_3 > AlCl_3$. Plants absorb a solution more actively if there are ions that do not interfere with metabolism, but apparently, more active absorption of a ‘bad’ aluminum ions also occurs.

Effects of aluminum toxicity. The root tip is deformed (Fig. 1, p. 213). The root becomes thickened; cell division continues but the stretching processes slow down. Our results show that growth in root tips is connected with the ability to